

THE BRIGHTEST OH MASER IN THE SKY: A FLARE OF EMISSION IN W75 N.

A.V. Alakoz,^{1,*} V.I. Slysh,¹ M.V. Popov,¹ and I.E. Val'tts¹

¹*Astro Space Center of the Lebedev Physical Center of RAS,
Profsoyuznaya Str. 84/32, 117997 Moscow, Russia*

A flare of maser radio emission in the OH-line 1665 MHz has been discovered in the star forming region W75 N in 2003, with the flux density of about 1000 Jy. At the time it was the strongest OH maser detected during the whole history of observations since the discovery of cosmic masers in 1965. The flare emission is linearly polarized with a degree of polarization near 100%. A weaker flare with a flux of 145 Jy was observed in this source in 2000 – 2001, which was probably a precursor of the powerful flare. Intensity of two other spectral features has decreased after beginning of the flare. Such variation of the intensity of maser condensation emission (increasing of one and decreasing of the other) can be explained by passing of the magneto hydrodynamic shock across regions of enhanced gas concentration.

Key words: interstellar medium, OH masers, variability.

* Electronic address: `rett@tanatos.asc.rssi.ru`

INTRODUCTION

W75 N is one of the active star formation regions in the Galaxy which is accompanied by the maser emission in lines of OH, methanol and H_2O . Mapping observations of the masers with high angular resolution has shown that the maser emission sources are connected with ultracompact HII regions (5; 9; 13), and possibly are located in protoplanetary disk surrounding massive stars. In this paper discovery of the powerful flare of emission is reported by which W75 N became the brightest OH cosmic maser for the whole history of observations.

OBSERVATIONS

Polarization observations of the maser emission from W75 N in all four Stokes parameters were conducted at the Nançay radio telescope (France) (4) in October 2003, and in two circular polarizations at the 64-m radio telescope in Kalyazin (Russia) (12) in July and October 2004, in OH lines 1665 and 1667 MHz. Digital autocorrelators with spectral resolution 0.137 km/s were used at both telescopes. The number of channels at Nançay was 8192, divided in 8 sections by 1024 channels each (both frequencies were observed, and each section corresponded to a separate polarization), the bandwidth was 0.781 MHz. At Kalyazin there was only one section of 4096 channels, with the bandwidth 3.125 MHz, so the measurements of the right hand and left hand circular polarizations had to be made sequentially. Additionally several records on the magnetic tape were made in the VLBI system S-2 on April 12, 2001 at 64-m radio telescope in Bear Lakes (Russia) and July 22, 2004 at Kalyazin. (The bandwidth in both cases was 8 MHz). The spectral analysis was performed with a software correlator, which allowed us to vary resolution in a wide range. The sensitivity of the Nançay radio telescope was 1.0 K/Jy, and that of Bear Lakes and Kalyazin radio telescopes was 0.65 K/Jy. The flux density calibration of Bear Lake and Kalyazin radio telescopes was performed with continuum sources Cyg A and 3C 274. The beam width of the Nançay radio telescope was $3'.5$ in the East-West direction and $19'$ in North-South direction. Bear Lakes and Kalyazin radio telescopes had circular beams with the width $8'$. In addition we have analyzed also observational data of this source in OH lines from NRAO VLBA archive for November 2000 and January 2001 (observations by Fish and Migenes).

RESULTS

Spectra of W75 N obtained with the VLBA in 1998 (13) and in 2001 (VLBA archive), at Nançay in 2003, and in Kalyazin in 2004, in the OH line 1665 MHz in right hand circular polarization are presented on Fig. 1. Some differences of spectra obtained with the VLBA from the rest three spectra arise because they were obtained in the VLBI regime while the single dish spectra were obtained in the total power regime. We do not show total power spectra from individual VLBA antennae, because they are corrupted by the contribution from nearby maser W75 S due to a large beam of VLBA antennae. In the first spectrum of 1998 which was obtained during VLBA mapping several features were observed which were marked as A,C,E,F,G,J and K in 13. They were present also in earlier observations of 1983 (3), approximately with the same intensity except features J and K at radial velocities 3.0 km/s and 0.65 km/s which had variable intensity. Subsequent VLBA observations in November 2000 and January 2001 (VLBA archive) have shown that the spectrum of the source in the radial velocity range from 4 km/s to 14 km/s, remained unchanged, while at lower radial velocities significant changes took place. A strong spectral feature at the radial velocity 2.3 km/s and with a flux density 145 Jy (75 Jy in the right hand circular polarization which is shown in Fig. 1) appeared, with the line width 0.35 km/s as well as a weaker feature at the radial velocity -0.2 km/s with a flux density 35 Jy. We marked them as P_1 and P_2 , respectively. At the same time the features J and K have diminished by a factor from 2 to 3. Subsequent observations in April 2001 with a 64-m telescope at Bear Lakes (spectrum is not shown) demonstrated that the flared features P_1 and P_2 dimmed by a factor from 2 to 5, remaining at the same radio velocities. One can suppose that the flare of the features P_1 and P_2 was a precursor of a still brighter flare of the maser emission which has been first observed by us 2.5 years later, on October 24, 2003 with the Nançay radio telescope. At the spectrum on Fig. 1 one can see a very bright feature at the radial velocity 1.8 km/c and the flux density 750 Jy and line width 0.35 km/s (N_1) as well as two weaker features N_2 at the radial velocity 0 km/s and N_3 at -1 km/s. Polarization measurements at Nançay show that all the features of the flare N_1 , N_2 , N_3 have high degree of linear polarization exceeding ($\geq 80\%$). The difference between radial velocities of the new features N_1 , N_2 and those of the precursor which amounts to about -0.5 km/s, and the decrease of the precursor in April 2001 mean that a flare of new spectral features was observed rather

than the brightening of old ones. At the same time the features from A to G have the same flux density and radial velocity. The features J and K, even if present, have the flux density less than 10 Jy. Subsequent observations conducted at Kalyazin in July 2004 confirmed the existence of the flared features practically at the same radial velocities and with the same flux densities. Later observations at Kalyazin in October-December 2004 reveal some decay of the strongest spectral feature N_1 accompanied by an increase of the weaker flare features N_2 and N_3 . In the second OH line at the frequency 1667 MHz, observed at Nançay, new spectral features have appeared also, but the intensity did not exceed 20 Jy and was comparable with the intensity of the rest of spectral feature.

DISCUSSION

Although the variability of OH masers is a well established phenomenon, there was no reports about OH maser emission flares which such a large flux density. Moreover, at present there is no known OH maser with a flux density equal or exceeding that of W75 N. A powerful flare of maser emission has been observed twice in the line of water vapor at 1.35 cm in Orion (1; 8; 10) with the flux above 10 MJy. Since the flux of water masers exceeds the flux of OH masers by a factor of $10 - 10^3$, there is no surprise that the scale of the flare is respectively larger for water maser.

In the paper of Slysh et al. (2002) a model of W75 N maser was proposed, in which the maser spots are located in disks rotating around two massive stars which in turn excite ultracompact HII-regions VLA 1 and VLA 2 (14) (Fig. 2). Spectral features from A to I at the radial velocities from 12.45 km/s to 3.70 km/s are connected with VLA 1 (some weaker features are not visible on the scale of our diagram and have no marks); only two features are connected with VLA 2: J and K at the radial velocities 3.0 km/s and 0.65 km/s, respectively. Judging on radial velocities of new spectral features of the flare 2 km/s and 0 km/s, they could be also connected with the VLA 2. VLA 1 and VLA 2 are separate centers of maser emission also in lines of H_2O and, probably, of methanol. These ultracompact HII regions are excited by young massive stars of spectral type B1 or B2 with a mass $10 M_\odot$ (11). The separation between the stars is about 1400 AU. In the paper of Torrelles et al. (14) a model of the source W75 N was proposed, in which the stars are sources of a wind and a shock which excite the maser emission. Maser system connected with VLA 2 is more compact

and, apparently, more active. The evidence of this comes from observations of high velocity masers in the OH 1667 MHz line in 1986 (6) and of the variability of H₂O maser emission with a period of 11.5 years, which are associated with VLA 2 (7). The new flare in the OH 1665 MHz line with the flux 1000 Jy, which is reported in this paper has occurred also in the maser system connected with the VLA 2. In the framework of the model of stellar wind shock one can assume that the flare began upon arrival of the shock into a region of enhanced abundance of H₂O and hydroxyl molecules and elevation of the kinetic temperature in this region, or was triggered by the enhancement of abundance of H₂O and hydroxyl molecules caused by the shock.

Almost simultaneously with the rising of the flux of spectral features of the precursor flare spectral features at the radial velocity 2.3 km/s and -0.2 km/s, a drop of the flux from spectral features J and K at the radial velocities 3.0 km/s and 0.65 km/s has occurred. One can see from Fig. 1 that these features are not visible in the spectrum of W75 N of 2001, at least at the level of 10 Jy, while in 1998 they had flux comparable with that of the brightest feature A (13). The features J and K were brighter in the past: the maximum flux of feature J was observed in 1993 (6); the feature was weaker in the earlier epochs of 1983 and 1986 (3; 6), as well as in later epoch 1998 (2; 13). Using this data one can estimate that the duration of feature J emission was $t=11$ years at the half intensity level, with the maximum in 1993. It is known from the VLBA maps (Fig. 2) (13) that the feature J as well as feature K is extended and has a shape of a thin strip 10.7×2.4 milliarcsec in extent, with the position angle 114° . At the distance to W75 N 2 kpc this corresponds to the linear size of the maser condensation $l_1 \times l_2 = (21.4 \times 4.8)$ AU. If one assumes that the emission from the feature J was caused by passing of a shock across the clump of matter with the above mentioned size, the velocity of the shock must be not less than $l_1/t = 9.3$ km/s, if the clump is in the plane of the sky, or more with an account of the projection effect.

Normal supersonic shocks are hardly able to propagate with such a velocity, since the sound velocity in molecular clouds is small, less than 1 km/s. However magneto hydrodynamic waves propagating with the Alfvén velocity may well be the agent exciting maser emission. It is known that in the maser emission regions magnetic field may exceed 5×10^{-3} Gauss, as follows from measurements of the Zeeman splitting. For example in W75 N two maser condensations have magnetic field 5.2×10^{-3} Gauss and 7.7×10^{-3} Gauss, respectively ([13]). Assuming the value of magnetic field equal to $H = 3 \times 10^{-3}$ Gauss and gas density $n_{H_2} = 10^5 -$

10^6 cm^{-3} , one gets Alfven velocity 15 km/s and 5 km/s, respectively, in agreement with the estimates of the excitation propagation velocity which ignited the maser emission of component J. One can suppose that after passing through the component J the excitation with the Alfven velocity has arrived to the next condensation which gave start to the next flare of the maser emission. Its intensity this time happened to be about 50 times higher than the intensity of component J; this can be explained by a more favorable conditions for the maser emission. More detailed information about temporal evolution of the maser emission flare can be obtained from results of high angular resolution VLBI mapping and from the space interferometer "Radioastron".

CONCLUSION

The brightest flare of emission in OH line since the beginning of observations in 1965 was discovered. The flare is connected with the activity of a young massive star which excites ultracompact HII region VLA 2 in the star forming region W75 N. The onset of the flare was accompanied by attenuation of emission from other maser condensations connected with the VLA 2. Such a behavior can be explained by propagation of magneto hydrodynamic shock through a region of enhanced gas concentration.

ACKNOWLEDGMENTS

Authors are grateful to collaborators from Astro Space Center of the Lebedev Physical Institute B.Z. Kanevskii, A.I. Smirnov, S.V. Logvinenko, V.I. Vasil'kov, S.V. Kalenskii, and also to Ray Escoffier (NRAO, USA) for the development of the receiver and correlator, and to Jean-Michel Martin, Eric Gerard (Meudon Observatory, France), Victor Migenes (University of Guanajuato, Mexico) for the help with observations, and to data analysts of NRAO for the access to the archive of results of observations.

This research was conducted under a partial support from the Russian Foundation for Basic Research (project code N 04-02-17057), CRDF (grant N RP1-2392-MO-02), Program for Fundamental Research of the Russian Academy of Science "Extended Objects in the Universe", Federal Program "Astronomy", and also in the framework of the preparation the observational program for the space interferometer "Radioastron".

-
1. Z. Abraham, N. L. Cohen, R. Opher, *et al.*, *Astron. Astrophys.* **100**, L10 (1981).
 2. A. L. Argon, M. J. Reid, and K. M. Menten, *Astrophys. J. Suppl. Ser.* **129**, 159 (2000).
 3. E. E. Baart, R. J. Cohen, R. D. Davies *et al.*, *MNRAS* **219**, 145 (1986).
 4. W. van Driel, J. Pezzani, and E. Gerard, *High Sensitivity Radio Astronomy* (Ed. N. Jackson, and R.J. Davis, Cambridge: Cambridge Univ. Press, 1996).
 5. T. R. Hunter, G. B. Taylor, M. Felli, *et al.*, *Astron. Astrophys.* **284**, 215 (1994).
 6. B. Hutawarakorn, R. J. Cohen, and G. C. Brebner, *MNRAS* **330**, 349 (2002).
 7. E. E. Lekht and V. V. Krasnov, *Astronomy Letters* **26**, 38 (2000).
 8. L. I. Matveyenko, *Sov. Astr. Letters*, **7**, 100 (1981).
 9. V. Minier, R. S. Booth, and J. E. Conway, *Astron. Astrophys.* **362**, 1093 (2000).
 10. T. Omodaka, T. Maeda, and N. Mochiduki, *IAU Circ. N* 6893 (1998).
 11. D. S. Shepherd, S. E. Kurtz, and L. Testi, *Astrophys. J.* **601**, 952 (2004).
 12. V. I. Slysh, M.V. Popov, B. Z. Kanevsky, *et al.*, *Astronomy Letters* **27**, 277 (2001).
 13. V. I. Slysh, V. Migenes, I. E. Val'tts, *et al.*, *Astrophys. J.* **564**, 317 (2002).
 14. J. M. Torrelles, J. F. Gómez, L. F. Rodríguez *et al.*, *Astrophys. J.* **489**, 744 (1997).




Figure 1. Spectra of OH maser W75 N in 1665 MHz line (right hand circular polarization), obtained at different epochs. 1998/VII: VLBA array, July 1, 1998; 2000/XI: VLBA array, November 22, 2000; 2003/X: Nancay radio telescope, October 24, 2003; 2004/VII: Kalyazin radio telescope, July 14, 2004; 2004/X: Kalyazin radio telescope, October 20, 2004; Spectral features marked by letters A — K, were identified in the paper of Slysh et al. (2002), features marked by letters P₁, P₂ and N₁, N₂ N₃ are new spectral features of the precursor (P) and flare (N). VLBA spectra are shown in interferometric mode, in which contribution from the nearby maser W75 S at the distance 14' is canceled.




Figure 2. Position of precursors P_1 and P_2 (in the vicinity of VLA 2) and the direction of the magnetic field in them (determined from VLBA archive observations in November 2000 and January 2001, Slysh, private communication). Features P_1 and P_2 are plotted on the VLBA map of OH maser condensations in W75 N, taken from Slysh et al. (2002). Circles show 1665 MHz features, triangles show 1667 MHz features. The direction of the magnetic field is shown by the vectors. The position of the ultracompact HII regions VLA 1 and VLA 2 are shown by an ellipse and a large circle, respectively. Dashed line shows a disk possibly connected with VLA 1.